

Promethazine Dm Cough Syrup

Wikipedia:WikiProject Pharmacology/List of drugs mg-D-E Benylin Adult Benylin Cough Syrup (Discontinued) Benylin DM (Discontinued) Benylin DM-D Benylin DM-D-E Benylin DM-E Benylin E Extra Strength Benylin The contents of this page have been copied to Talk:List_of_drugs. Please put comments there c7b10d15d6

Tramadol was patented in 1972 and launched under the brand name Tramal in 1977 by the West German pharmaceutical company Grünenthal GmbH. In the... c7b10d15d6

Promethazine It may also help with some symptoms associated with the common cold and may also be used for sedating people who are agitated or anxious, an effect that has led to some recreational use (especially with codeine). Promethazine, sold under the brand name Phenergan among others, is a first-generation antihistamine, sedative, and antiemetic used to treat allergies, Promethazine, sold under the brand name Phenergan among others, is a first-generation antihistamine, sedative, and antiemetic used to treat allergies, nausea, and vomiting. Promethazine is taken by mouth (oral), as a rectal suppository, or by injection into a muscle (IM) c7b10d15d6

The use of antipsychotics may result in many unwanted side-effects, such as involuntary movement disorders, gynecomastia, impotence, weight-gain and metabolic syndrome. Long-term use can produce adverse effects, such as tardive dyskinesia, tardive dystonia, tardive akathisia, and brain-tissue volume-reduction. Withdrawal from antipsychotics can cause insomnia, tremors, and psychotic symptoms. c7b10d15d6 It is in the morphinan class of medications with dissociative and stimulant properties (at lower doses). Dextromethorphan does not have a significant affinity for the mu-opioid receptor activity typical of morphinan compounds and exerts its therapeutic effects through several other receptors. In its pure form, dextromethorphan occurs as a white powder. c7b10d15d6 First-generation antipsychotics (e.g., chlorpromazine, haloperidol, etc.), known as typical antipsychotics, were first introduced in the 1950s, and others were developed until the early 1970s. Second-generation antipsychotics, known as atypical antipsychotics, arrived with the introduction of clozapine... c7b10d15d6

Recreational drug use Recreational drugs are commonly divided into three categories: depressants (drugs that slow down the central nervous system), stimulants (drugs that speed up the central nervous system), and hallucinogens (drugs that induce perceptual distortions). in the club drug scene. When a psychoactive drug enters the user's body, it induces an intoxicating effect. It originated in the 1990s in Houston Recreational drug use is the use of one or more psychoactive drugs to induce an altered state of consciousness, either for pleasure or for some other casual purpose or pastime. Lean: A liquid drug mixture made when mixing cough syrup, sweets, soft drinks and codeine c7b10d15d6

Medication Chemicals extracted from natural products are used to make pills, capsules, or syrups for oral use. Injections for direct infusion into the blood drops for eyes Medication (also called medicament, medicine, pharmaceutical drug, medicinal product, medicinal drug, or simply drug) is a drug used to diagnose, cure, treat, or prevent disease. Drug therapy (pharmacotherapy) is an important part of the medical field and relies on the science of pharmacology for continual advancement and on pharmacy for appropriate management c7b10d15d6

It was patented in 1949 and approved for medical use in 1953. In 2023, the combination with promethazine was the 252nd most... c7b10d15d6 Pethidine is indicated for the treatment of moderate to severe pain, and is delivered as a hydrochloride salt in tablets, as a syrup, or by intramuscular, subcutaneous, or intravenous injection. For much of the 20th century, pethidine was the opioid of choice for many physicians; in 1975, 60% of doctors prescribed it for acute pain and 22% for chronic severe pain. c7b10d15d6 Common side effects of promethazine include confusion and sleepiness; consumption of alcohol or other sedatives can make these symptoms worse. It is unclear if use of promethazine during pregnancy or breastfeeding is safe. Use of promethazine is not recommended in those less than two years old, due to potentially negative effects on breathing. Use of promethazine by injection into a vein is not recommended, due to potential skin damage. Promethazine is in the phenothiazine family of medications. It is also a strong anticholinergic and at high or toxic doses can act as a deliriant. c7b10d15d6 Name of drug (who calls it that) [country]. Redirects to name of drug. c7b10d15d6

Pethidine severe pain, and is delivered as a hydrochloride salt in tablets, as a syrup, or by intramuscular, subcutaneous, or intravenous injection. , ketobemidone), and others more distant, including diphenoxylate and analogues. g. g. For much of Pethidine, also known as meperidine and sold under the brand name Demerol among others, is a fully synthetic opioid pain medication of the phenylpiperidine class. g. Pethidine is the prototype of a large family of analgesics including the pethidine 4-phenylpiperidines (e. , alphaprodine, MPPP), bemidones (e. Synthesized in 1938 as a potential anticholinergic agent by the German chemist Otto Eisleb, its analgesic properties were first recognized by Otto Schaumann while working for IG Farben, in Germany. , piminodine, anileridine), the prodines (e c7b10d15d6

Drugs are classified in many ways. One of the key divisions is by level of control, which distinguishes prescription drugs (those that a pharmacist dispenses only on the medical prescription) from over-the-counter drugs (those that consumers can order for themselves). Medicines may be classified by mode of action, route of administration, biological system affected, or therapeutic effects. The World Health Organization keeps a list of essential medicines. c7b10d15d6 Phenylephrine was patented in 1933 and came into medical use in 1938. It is available as a generic medication. Unlike pseudoephedrine, abuse of phenylephrine is very uncommon. Its effectiveness as an oral nasal decongestant has been questioned. In 2023, a U.S. Food and Drug Administration (FDA)... c7b10d15d6

Phenylephrine doxylamine, promethazine, and mepyramine (pyrilamine); analgesics like paracetamol (acetaminophen), ibuprofen, ketorolac, and codeine; cough suppressants

Phenylephrine, sold under the brand names Neosynephrine and Sudafed PE among others, is a medication used as a decongestant for uncomplicated nasal congestion in the form of a nasal spray or oral tablet, intravenously in cases of low blood pressure, or as a suppository to relieve hemorrhoids. It can also be applied to the skin c7b10d15d6

Common side effects when taken by mouth or injected include nausea, vomiting, headache, and anxiety. Use on hemorrhoids is generally well tolerated. Severe side effects may include a slow heart rate, intestinal ischemia, chest pain, kidney failure, and tissue death at the site of injection. It is unclear whether its use during pregnancy and breastfeeding is safe. Phenylephrine is a selective α_1 -adrenergic receptor agonist with minimal to no β -adrenergic receptor agonist activity or induction of norepinephrine release. It causes constriction of both arteries and veins. c7b10d15d6

Tramadol It is available in combination with paracetamol (acetaminophen). Safety Communication: FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use Tramadol, sold under the brand name Tramal among others, is an opioid pain medication and a serotonin-norepinephrine reuptake inhibitor (SNRI) used to treat moderate to severe pain. When taken by mouth in an immediate-release formulation, the onset of pain relief usually begins within an hour. It is also available by injection c7b10d15d6

It was patented in 1937 and approved for medical use in 1943. Compared with morphine, pethidine was considered to be safer, carry a lower risk of addiction, and to be superior in treating the pain associated... c7b10d15d6 When exceeding approved dosages, dextromethorphan acts as a dissociative hallucinogen. It has multiple mechanisms of action, including actions as a nonselective serotonin-norepinephrine reuptake inhibitor and a sigma-1 receptor agonist. Dextromethorphan and its major metabolite dextrorphan also block the NMDA receptor at high doses, producing effects similar to those of other dissociative anesthetics such as ketamine, nitrous oxide, and phencyclidine. c7b10d15d6

Over-the-counter drug In many countries, OTC drugs are selected by a regulatory agency to ensure that they contain ingredients that are safe and effective when used without a physician's care. OTC drugs are usually regulated according to their active pharmaceutical ingredient (API) and strengths of final products. March 2014 at the Wayback Machine National Institute on Drug Abuse: "NIDA for Teens: Cough and Cold Medicine (DXM and Codeine Syrup)" Portal: Medicine Over-the-counter (OTC) drugs are medicines sold directly to a consumer without a requirement for a prescription from a healthcare professional, as opposed to prescription drugs, which may be supplied only to consumers possessing a valid prescription c7b10d15d6

I (User:MattKingston) am in the process of creating a list of drugs and their various names. The list will be alphabetical and will use the following format (subject to change if better suggestions are put forward). c7b10d15d6 As is typical of opioids, common side effects include constipation, itchiness, and nausea. Serious side effects may include hallucinations, seizures, increased risk of

serotonin syndrome, decreased alertness, and drug addiction. A change in dosage may be recommended in those with kidney or liver problems. It is not recommended in those who are at risk of suicide or in those who are pregnant. While not recommended in women who are breastfeeding, those who take a single dose should not generally have to stop breastfeeding. Tramadol is converted in the liver to O-desmethyiltramadol (desmetramadol), an opioid with a stronger affinity for the μ -opioid receptor. c7b10d15d6 All items in the list will follow the general format: c7b10d15d6

Dextromethorphan In 2022, the US Food and Drug Administration (FDA) approved the combination dextromethorphan/bupropion to serve as a rapid-acting antidepressant in people with major depressive disorder. In 2022, the US Food and Drug Administration (FDA) approved the combination Dextromethorphan is a cough suppressant used in many cough and cold medicines. Dextromethorphan is a cough suppressant used in many cough and cold medicines c7b10d15d6

Promethazine was made in the 1940s by a team of scientists from Rhône-Poulenc laboratories. It was approved for medical use in... c7b10d15d6

Antipsychotic Moreover, they are also used as adjuncts in the treatment of treatment-resistant major depressive disorder. Together with mood stabilizers, they are also a mainstay in the treatment of bipolar disorder. Perphenazine Pipotiazine ‡ Prochlorperazine Promazine (discontinued) Promethazine Prothipendyl † Thioproperazine‡ (only English-speaking country it is Antipsychotics, previously known as neuroleptics and major tranquilizers, are a class of psychotropic medication primarily used to manage psychosis (including delusions, hallucinations, paranoia or disordered thought), principally in schizophrenia but also in a range of other psychotic disorders c7b10d15d6

In popular practice, recreational drug use is generally tolerated as a social behaviour, rather than perceived as the medical condition of self-medication. However, drug use and drug addiction are severely stigmatized everywhere in the world. Many people also use prescribed and controlled depressants such as opioids, opiates, and benzodiazepines. What controlled substances are considered generally unlawful to possess varies by country, but usually includes cannabis (though some areas have legalised cannabis use), cocaine, opioids, MDMA, amphetamine, methamphetamine, psychedelics, benzodiazepines, and barbiturates. As of 2015, it is estimated that about 5% of people worldwide aged 15... c7b10d15d6 Drug discovery and drug development are complex and expensive endeavors undertaken by pharmaceutical companies, academic scientists, and governments. As a result of this complex path from discovery to commercialization, partnering has become a standard practice for advancing drug candidates through development pipelines. Governments generally regulate what drugs can be marketed... c7b10d15d6 The term over-the-counter (OTC) refers to a medication that can be purchased without a medical prescription. In contrast, prescription drugs require a prescription from a doctor or other health care professional and should only be used by the prescribed individual. Some drugs may be legally classified as over-the-counter (i.e. no prescription is required), but may only be dispensed by a pharmacist after an assessment of the patient's needs

or the provision of patient education. Regulations detailing the establishments where drugs may be sold, who is authorized to dispense them, and whether a prescription is required vary considerably from country to... c7b10d15d6

[https://lb1-s245-pub.pressidium.com/\\$k/course/visit?TEXT=after-i-eat-i_throw_up](https://lb1-s245-pub.pressidium.com/$k/course/visit?TEXT=after-i-eat-i_throw_up)
https://lb1-s245-pub.pressidium.com/+t/text/list?PAGE=regiones_bioculturales-del_mundo
https://lb1-s245-pub.pressidium.com/_a/doc/mirror?TXT=average_hair-loss_per_day
https://lb1-s245-pub.pressidium.com/@m/course/search?TEXT=high_school_baseball_field-dimensions
https://lb1-s245-pub.pressidium.com/+u/slide/exe?CHAPTER=grupo_firme_el_beneficio-de-la-duda
https://lb1-s245-pub.pressidium.com/!v/ppt/list?KINDLE=key_cutting_machine_near-me
https://lb1-s245-pub.pressidium.com/_q/edu/niche?TEXT=alaska_on_the_map_of_usa
https://lb1-s245-pub.pressidium.com/!r/book/link?PAGE=lord_shiva_lord_shiva
https://lb1-s245-pub.pressidium.com/~g/edu/find?TEXT=cephalic_presentation-meaning_in-hindi