

Side Effects Of Effexor Antidepressant

Abilify (aripiprazole) - atypical antipsychotic used to treat schizophrenia, bipolar disorder, and irritability associated with autism 331893f79c SSRIs are the most widely prescribed antidepressants in many countries. In adults, they are recommended as a first-line treatment for moderate to severe depression, while for mild depression non-drug treatments are preferred unless the patient chooses medication. A 2018 meta analysis of 21 antidepressants showed that all antidepressants (including SSRIs) were more effective than placebo in adults with major depressive... 331893f79c Though psychiatric medication is the most frequently prescribed therapy for major depression, psychotherapy may be effective, either alone or in combination with medication. Given an accurate diagnosis of major depressive disorder, in general the type of treatment (psychotherapy or antidepressants, alternate or other treatments, or active intervention) is "less important than getting depressed patients involved in an active therapeutic program." 331893f79c Discontinuation syndrome was coined by Eli Lilly as a way to distance antidepressants from other dependence-forming psychotropics with higher addiction risk profiles. However, the evidence supports that what has been mislabeled as discontinuation... 331893f79c Insomnia 331893f79c SSRIs primarily work by blocking serotonin reabsorption (reuptake) via the serotonin transporter, leading to gradual changes in brain signaling and receptor regulation, with some also interacting with sigma-1 receptors, particularly fluvoxamine, which may contribute to cognitive effects. Marketed SSRIs include six main antidepressants—citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, and sertraline—and dapoxetine, which is indicated for premature ejaculation. Fluoxetine has been approved for veterinary use in the treatment of canine separation anxiety. 331893f79c Common side effects of antidepressants include dry mouth, weight gain, dizziness, headaches, akathisia, sexual dysfunction, and emotional blunting. There is an increased risk of suicidal thinking and behavior when taken by children, adolescents, and young adults. Discontinuation syndrome, which resembles recurrent depression in the case of the SSRI class, may occur after stopping the intake of any antidepressant, having effects which may be permanent and irreversible. Tapering off medications gradually is shown to reduce the risk of withdrawal complications. 331893f79c Anxiety is a naturally occurring emotion and response. When anxiety levels exceed the tolerability of a person, anxiety disorders may occur. People with anxiety disorders can exhibit fear responses, such as defensive behaviors, high levels of alertness, and negative emotions. Those with anxiety disorders may have concurrent psychological disorders, such as depression. Anxiety disorders are classified using six possible clinical assessments: 331893f79c Healy has been involved as an expert witness in homicide and suicide trials involving psychotropic drugs, and has brought concerns about some medications to the attention of drug regulators. He has also said that pharmaceutical companies sell drugs by marketing diseases and co-opting academic opinion-leaders. In his 2012 book Pharmageddon he argues that pharmaceutical companies have dominated healthcare in America, often with life-threatening results for patients. Healy is a founder and chief executive officer of Data Based Medicine Limited, which aims to make medicines safer through... 331893f79c Psychotherapy is the treatment of choice in those under the age of 18, with medication offered only in conjunction with the former and generally not as a first line agent. The possibility of depression, substance misuse or other mental health problems in the... 331893f79c Nausea — an adverse effect that is more common with venlafaxine than with the SSRIs. Usually transient and less severe in those receiving the extended release formulations. 331893f79c Ejaculation disorder — sexual side effects can be seen with virtually any antidepressant, especially those that inhibit the reuptake of serotonin (including venlafaxine). 331893f79c

List of psychotropic medications narcolepsy Effexor and Effexor XR (venlafaxine) - an antidepressant of the SNRI class Elavil (amitriptyline) - a tricyclic antidepressant used as a first-line This is a list of psychotropic medications that are currently being marketed, alphabetized by brand name 331893f79c

Adderall (mixed amphetamine salts) - a stimulant used to treat ADHD 331893f79c == A == 331893f79c Dizziness 331893f79c Headache — an often transient side effect that is common to most serotonin reuptake inhibitors and that most often occurs at the beginning of therapy or after a dose escalation. 331893f79c The human serotonin transporter (SERT) and norepinephrine transporter (NET) are membrane transport proteins that are responsible for the reuptake of serotonin and norepinephrine from the synaptic cleft back into the presynaptic nerve terminal. Dual inhibition of serotonin and norepinephrine reuptake can offer advantages over... 331893f79c Ambien (zolpidem) - nonbenzodiazepine used as a sleep aid 331893f79c Depression is a symptom of some physical diseases; a side effect of some drugs and medical treatments; and a symptom of some mood disorders such as major depressive disorder or dysthymia. Physical causes are ruled out with a clinical assessment of depression that measures vitamins, minerals, electrolytes, and hormones. 331893f79c Aricept (donepezil) - used to slow the progression of Alzheimer's disease 331893f79c

Antidepressant disorder, anxiety disorders, chronic pain, and addiction. Common side effects of antidepressants include dry mouth, weight gain, dizziness, headaches, akathisia Antidepressants, also known in the past as psychic energizers, are a class of medications used to treat major depressive disorder, anxiety disorders, chronic pain, and addiction 331893f79c

== Very common (>10% incidence) == 331893f79c Updated May 2026 (List is Complete) 331893f79c

Selective serotonin reuptake inhibitor reuptake inhibitors (SSRIs) are a class of drugs that are typically used as antidepressants in the treatment of major depressive disorder, anxiety disorders Selective serotonin reuptake inhibitors (SSRIs) are a class of drugs that are typically used as antidepressants in the treatment of major depressive disorder, anxiety disorders, and other psychiatric conditions 331893f79c

Antabuse (disulfiram) - inhibits the enzyme acetaldehyde dehydrogenase, causing acetaldehyde poisoning when ethanol is consumed; used to cause severe hangover when drinking; increases liver, kidney, and brain

damage from drinking 331893f79c Anquil (benperidol) – an antipsychotic primarily used to control antisocial hypersexual behaviour 331893f79c Common side effects include loss of appetite, constipation, dry mouth, dizziness, sweating, insomnia, drowsiness and sexual problems. In some patients, sexual dysfunction may persist even after the drug is discontinued, a condition known as post-SSRI sexual dysfunction. Severe side effects include an increased risk of suicide, mania, and serotonin syndrome. Antidepressant withdrawal syndrome may occur if stopped. A meta-analysis of randomized trials in depression found an increased rate of serious adverse events, particularly sexual dysfunction and anorexia... 331893f79c

Venlafaxine It is used to treat major depressive disorder, generalized anxiety disorder, panic disorder, and social anxiety disorder. It is also available as the salt venlafaxine besylate (venlafaxine benzenesulfonate monohydrate) in an extended-release formulation (Venbysi XR). Studies have shown that venlafaxine improves post-traumatic stress disorder (PTSD) as a recommended first-line treatment. It may also be used for chronic neuropathic pain. Venlafaxine, sold under the brand name Effexor among others, is an antidepressant medication of the serotonin–norepinephrine reuptake inhibitor (SNRI) Venlafaxine, sold under the brand name Effexor among others, is an antidepressant medication of the serotonin–norepinephrine reuptake inhibitor (SNRI) class. It is taken orally (swallowed by mouth) 331893f79c

Aponal, Quitaxon, Sinequan (doxepin) – a tricyclic antidepressant used to treat nerve pain, insomnia; similar to imipramine 331893f79c Ativan (lorazepam) – a benzodiazepine, used to treat anxiety 331893f79c Asthenia (weakness) 331893f79c

David Healy (psychiatrist) contribution of antidepressants to suicide, conflict of interest between pharmaceutical companies and academic medicine, and the history of pharmacology David Healy FRCPsych, a professor of psychiatry at Bangor University in the United Kingdom, is a psychiatrist, psychopharmacologist, scientist and author. Healy has written more than 150 peer-reviewed articles, 200 other articles, and 20 books, including The Antidepressant Era, The Creation of Psychopharmacology, The Psychopharmacologists Volumes 1–3, Let Them Eat Prozac and Mania: A Short History of Bipolar Disorder. His main areas of research are the contribution of antidepressants to suicide, conflict of interest between pharmaceutical companies and academic medicine, and the history of pharmacology 331893f79c

Management of depression Venlafaxine (Effexor) from the SNRI class Management of depression is the treatment of depression that may involve a number of different therapies: medications, behavior therapy, psychotherapy, and medical devices. inhibitors can be used as antidepressants. Norepinephrine reuptake inhibitors (NRIs) can be used as antidepressants 331893f79c

Asendin (amoxapine) – an dibenzoxazepine antidepressant 331893f79c The effectiveness of antidepressants for treating depression in adults has strong support, though studies also highlight potential risks and limitations. In children and adolescents, evidence of efficacy is more limited, despite a marked increase in antidepressant prescriptions for these age groups since the 2000s. A 2018 meta-analysis reported that the 21 most commonly prescribed antidepressants were found in all studies to be more effective than placebos... 331893f79c

Serotonin–norepinephrine reuptake inhibitor SNRIs are monoamine reuptake inhibitors; specifically, they inhibit the reuptake of serotonin and norepinephrine. are relatively few of them. SNRIs can be contrasted with the selective serotonin reuptake inhibitors (SSRIs) and norepinephrine reuptake inhibitors (NRIs), which act upon single neurotransmitters. These neurotransmitters are thought to play an important role in mood regulation. However, the SNRIs are among the most widely used antidepressants today. Off-label uses include treatments for attention deficit hyperactivity disorder (ADHD) and obsessive–compulsive disorder (OCD). Since norepinephrine pathways play a role in pain modulation, SNRIs are also widely used for chronic pain conditions such as neuropathic pain and fibromyalgia. In 2009, Cymbalta and Effexor were the 11th- and Serotonin–norepinephrine reuptake inhibitors (SNRIs) are a class of antidepressant medications used to treat major depressive disorder (MDD), anxiety disorders, chronic neuropathic pain, fibromyalgia, and menopausal symptoms 331893f79c

Withdrawal 331893f79c Sweating 331893f79c

Anxiolytic venlafaxine (Effexor), entered the market in 1993. Anxiolytic medications are used for the treatment of anxiety disorders and their related psychological and physical symptoms. SNRIs can target serotonin and norepinephrine transporters while avoiding imposing significant effects on other An anxiolytic (; also antipanic or anti-anxiety agent) is a medication or other intervention that reduces anxiety. This effect is in contrast to anxiogenic agents which increase anxiety 331893f79c

Azstarys (Serdexmethylphenidate/Dexmethylphenidate) - a long-acting... 331893f79c

Antidepressant discontinuation syndrome Psychosis may rarely occur. If stopped too quickly, a withdrawal injury can occur. This is referred to as protracted withdrawal and may last for several months or years. The symptoms may include dizziness, vertigo, postural orthostatic tachycardia syndrome (POTS), tinnitus, insomnia, nausea, poor balance, sensory changes, "brain zaps", emotional lability or extreme emotional changes, rage, suicidal ideation, akathisia, dysesthesia, intrusive thoughts, depersonalization, and derealization, mania, anxiety, depression, stuttering, and flu-like symptoms. The symptoms may include dizziness Antidepressant discontinuation syndrome, or antidepressant withdrawal, is a condition that can occur following switching, reducing, or discontinuing an antidepressant medication following its continuous use of at least a month. Depending on the specific antidepressant's half-life, withdrawal can begin within a few days or weeks, but late onset or delayed onset withdrawal can occur months after cessation. following switching, reducing, or

discontinuing an antidepressant medication following its continuous use of at least a month 331893f79c

== Nature of anxiety == 331893f79c

List of adverse effects of venlafaxine Ejaculation disorder — sexual side effects can be seen with virtually any antidepressant, especially those that inhibit the reuptake of serotonin (including venlafaxine) The following list shows the rates of adverse symptoms seen in people taking venlafaxine 331893f79c

Anafranil (clomipramine) - a tricyclic antidepressant; mostly used to treat OCD 331893f79c Somnolence 331893f79c Dry mouth 331893f79c

[https://lb1-s245-pub.pressidium.com/\\$q/chap/goto?PAGE=brazilian-lymphatic_drainage_massage](https://lb1-s245-pub.pressidium.com/$q/chap/goto?PAGE=brazilian-lymphatic_drainage_massage)
https://lb1-s245-pub.pressidium.com/-g/ebook/file?EBOOK=how_to_find-population-mean
https://lb1-s245-pub.pressidium.com/=a/chap/data?PDF=headache-and_diarrhea_and_nausea
https://lb1-s245-pub.pressidium.com/_z/edu/list?EBOOK=1_1-4-pounds_in-grams
https://lb1-s245-pub.pressidium.com/_i/ppt/slug?PUB=problems_with_problem_based_learning
https://lb1-s245-pub.pressidium.com/@k/ref/go?EPUB=what_is-human_machine_interaction
[https://lb1-s245-pub.pressidium.com/\\$r/ref/search?TEXT=decimas-de_fiebre_sin_motivo-aparente_en_adultos](https://lb1-s245-pub.pressidium.com/$r/ref/search?TEXT=decimas-de_fiebre_sin_motivo-aparente_en_adultos)
https://lb1-s245-pub.pressidium.com/_m/short/url?CHAPTER=4-3_kilos_to_pounds