

Side Effects Of Dates In Pregnancy

First Trimester

It is on the World Health Organization's List of Essential Medicines. In some countries it is available as a generic medication. It was approved for medical use in the United States in May 2026. 402282fc91 Common side effects include vomiting, diarrhea, rash, and increased liver enzymes. Serious side effects may include liver problems, QT prolongation, and seizures. Use during pregnancy may increase the risk of miscarriage or birth defects. Fluconazole is in the azole antifungal family of medication. It is believed to work by affecting the fungal cellular membrane. 402282fc91

Miscarriage use. About 80% of miscarriages occur in the first 12 weeks of pregnancy (the first trimester). The term miscarriage is sometimes used to refer to all forms of pregnancy loss and pregnancy with abortive outcomes before 20 weeks of gestation. Miscarriage before six weeks of gestation is defined as biochemical loss by ESHRE. Once ultrasound or histological evidence shows that a pregnancy has existed, the term used is clinical miscarriage, which can be "early" (before 12 weeks) or "late" (between 12 and 21 weeks). The underlying cause in about half of cases involves chromosomal Miscarriage, also known in medical terms as a spontaneous abortion, is an end to pregnancy resulting in the loss and expulsion of an embryo or fetus from the womb before it can survive independently. Spontaneous fetal termination after 20 weeks of gestation is known as a stillbirth 402282fc91

Pregnancy Pregnancy may be confirmed with a pregnancy test. A multiple pregnancy involves more than one offspring, such as with twins. Pregnancy is divided into three trimesters of Pregnancy is the time during which one or more offspring gestates inside a woman's uterus. Methods of contraception are used to avoid pregnancy 402282fc91

Dolutegravir/lamivudine/tenofovir It is taken by mouth. While there are concerns that use during pregnancy results in a 0. As of 2019, it is listed by the World Health Organization (WHO) as the first line treatment for adults, with tenofovir/lamivudine/efavirenz as an alternative. by mouth. Side effects may include trouble sleeping, weight gain, and rash. It is a combination of dolutegravir, lamivudine, and tenofovir disoproxil. 2% increased Dolutegravir/lamivudine/tenofovir (DTG/3TC/TDF) is a fixed-dose combination antiretroviral medication used to treat HIV/AIDS 402282fc91

The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) recommends that pregnant women be offered routine obstetric ultrasounds between 18 weeks' and 24 weeks' gestational age (the anatomy scan) in order to confirm pregnancy dating, to measure the fetus so that growth abnormalities can be recognized quickly later in pregnancy, and to assess for congenital malformations

and multiple pregnancies (twins, etc.). Additionally, the ISUOG recommends that pregnant patients who desire genetic testing have obstetric ultrasounds between 11 weeks' and 13 weeks 6 days' gestational age in countries with resources to perform them (the nuchal scan). Performing an ultrasound at this early stage of pregnancy can more accurately confirm... 402282fc91 == Causes == 402282fc91 The most common symptom of a miscarriage is vaginal bleeding, with or without pain. Tissue and clot-like material may leave the uterus and pass through and out of the vagina. Risk factors for miscarriage include being an older parent, previous miscarriage, exposure to tobacco smoke, obesity, diabetes, thyroid problems, and drug or alcohol use. About 80% of miscarriages occur in the first 12 weeks of pregnancy (the first trimester). The underlying cause in about half of cases involves chromosomal... 402282fc91

Fluconazole It is given either by mouth or by injection into a vein. It is also used to prevent candidiasis in those who are at high risk such as following organ transplantation, low birth weight babies, and those with low blood neutrophil counts. torsades de pointes In 2011, the US FDA reports that treatment with chronic, high doses of fluconazole during the first trimester of pregnancy may be associated Fluconazole, sold under the brand name Diflucan among others, is an antifungal medication used for a number of fungal infections. These include candidiasis, blastomycosis, coccidioidomycosis, cryptococcosis, histoplasmosis, dermatophytosis, and tinea versicolor 402282fc91

Also, pregnancy can cause hypercoagulability by other factors, e... 402282fc91 Many of the early signs of pregnancy will be similar to symptoms that come right before a period, and it can be hard to tell the difference. 402282fc91

Hypercoagulability in pregnancy However, when combined with an additional underlying hypercoagulable states, the risk of thrombosis or embolism may become substantial. Pregnancy itself is a factor of hypercoagulability (pregnancy-induced hypercoagulability), as a physiologically adaptive mechanism to prevent post partum bleeding. Also, pregnancy can cause hypercoagulability Hypercoagulability in pregnancy is the propensity of pregnant women to develop thrombosis (blood clots). may occur at the end of the first trimester, due to enhanced compliance of the vessel walls by a hormonal effect 402282fc91

Drugs in pregnancy The Food and Drug Administration (FDA) in the United States reports that there are six million pregnancies with at least 50% of the women taking at least one medication. Some of the consequences on the babies include physical and mental abnormalities, higher risk of stillbirth, neonatal abstinence syndrome (NAS), sudden infant death syndrome (SIDS), low birthweight, and others. In addition a reported 5-10% of women of childbearing age use alcohol or other addictive substances. and recreational drugs, may have effects during pregnancy on the pregnant woman and fetus that vary from the effects of the drug on people who are not pregnant Drugs, including medications and recreational drugs, may have effects during pregnancy on the pregnant woman and fetus that vary from the effects of the drug on people who are not pregnant. Of those who bear children, recreational drug use can have serious consequences to the health of not only the mother, but

also the fetus as many medications can cross the placenta and reach the fetus
402282fc91

== Medical uses == 402282fc91

Beginning of pregnancy controversy Because an abortion is defined as ending an established pregnancy, rather than as destroying a fertilized egg, depending on when pregnancy is considered to begin, some methods of birth control as well as some methods of infertility treatment might be classified as causing abortions. Controversy over the beginning of pregnancy occurs in different contexts, particularly as it is discussed within the debate of abortion in the United States. Because Controversy over the beginning of pregnancy occurs in different contexts, particularly as it is discussed within the debate of abortion in the United States
402282fc91

Side effects may include trouble sleeping, weight gain, and rash. While there are concerns that use during pregnancy results in a 0.2% increased risk of neural tube defects in the baby, this does not rule out its use. Use remains recommended after the first trimester. Use is not recommended in those with kidney problems. The combination is a type of antiretroviral therapy. 402282fc91

Signs and symptoms of pregnancy bleeding in the first trimester or bleeding associated with pain, may be a sign of a complication, such as a miscarriage or an ectopic pregnancy, that would Signs and symptoms of pregnancy are common, benign conditions that result from the changes to the body that occur during pregnancy. If a person is concerned about their symptoms they should be encouraged to speak with an appropriate healthcare professional. Signs and symptoms of pregnancy typically change as pregnancy progresses, although several symptoms may be present throughout. Depending on severity, common symptoms in pregnancy can develop into complications. Each pregnancy can be quite different and many people do not experience the same or all of the symptoms. Pregnancy symptoms may be categorized based on trimester as well as region of the body affected 402282fc91

== Side effects and mechanism == 402282fc91 Conception usually occurs following vaginal intercourse, but can also occur through assisted reproductive technology procedures. A pregnancy may end in a live birth, a miscarriage, an induced abortion, or a stillbirth. Childbirth typically occurs around 40 weeks from the start of the last menstrual period (LMP), a span known as the gestational age; this is just over nine months. Counting by fertilization age, the length is about 38 weeks. Implantation occurs on average 8–9 days after fertilization. An embryo is the term for the developing offspring during the first seven weeks following implantation (i.e. ten weeks' gestational age), after which the term fetus is used until the birth of a baby. 402282fc91 Pregnancy is divided into three trimesters of approximately... 402282fc91 When done legally in industrialized societies, induced abortion is one of the safest procedures in medicine. Modern methods use medication or surgery for abortions. The drug mifepristone (aka RU-486) in combination with prostaglandin appears to be as safe and effective as surgery during the first and second trimesters

of pregnancy. Self-managed medication abortion is highly effective and safe throughout the first trimester. The most common surgical technique involves dilating the cervix and using... 402282fc91

Obstetric ultrasonography 1056/NEJMra1302417 Obstetric ultrasonography, or prenatal ultrasound, is the use of medical ultrasound in pregnancy, in which sound waves are used to create real-time visual images of the developing embryo or fetus in the uterus (womb). doi:10. New England Journal of Medicine. "Diagnostic Criteria for Nonviable Pregnancy Early in the First Trimester". The procedure is a standard part of prenatal care in many countries, as it can provide a variety of information about the health of the mother, the timing and progress of the pregnancy, and the health and development of the embryo or fetus. (ed. 369 (15): 1443-1451.) 402282fc91

Abortion Other reasons include maternal health, an inability to afford a child, domestic violence, lack of support, feelings of being too young, wishing to complete an education or advance a career, and not being able, or willing, to raise a child conceived as a result of rape or incest. Abortion occurring without intervention is known as spontaneous abortion or "miscarriage", and occurs in roughly 30-40% of all pregnancies. RU-486) in combination with prostaglandin appears to be as safe and effective as surgery during the first and second trimesters of pregnancy. Self-managed Abortion is the termination of a pregnancy by removal or expulsion of an embryo or fetus. The unmodified word abortion generally refers to induced abortion, or deliberate actions to end a pregnancy. Common reasons for inducing an abortion are birth-timing and limiting family size 402282fc91

The controversy is not primarily a scientific issue, since knowledge of human reproduction and development has become very refined; the linguistic questions remain debated for other reasons. The issue poses larger social, legal, medical, religious, philosophical, and political ramifications because some people, such as Concerned Women for America, identify the beginning of a pregnancy as the beginning of an individual human being's life. Many of these arguments are related to the anti-abortion movement. In this way of thinking, if the pregnancy has not yet begun, then stopping the process is not abortion and therefore can contain none of the moral issues associated with abortion, but if it is a pregnancy, then stopping it is a morally significant... 402282fc91 Pregnancy-induced hypercoagulability is probably a physiologically adaptive mechanism to prevent post partum hemorrhage. Pregnancy changes the plasma levels of many clotting factors, such as fibrinogen, which can rise up to three times its normal value. Thrombin levels increase. Protein S, an anticoagulant, decreases. However, the other major anticoagulants, protein C and antithrombin III, remain constant. Fibrinolysis is impaired by an increase in plasminogen activator inhibitor-1 (PAI-1 or PAI) and plasminogen activator inhibitor-2 (PAI-2), the latter synthesized from the placenta. Venous stasis may occur at the end of the first trimester, due to enhanced compliance of the vessel walls by a hormonal effect. 402282fc91 As of 2019, it is listed by the World Health Organization (WHO) as the first-line treatment for adults with HIV/AIDS, with tenofovir/lamivudine/efavirenz as an alternative. It may be used in people with both

HIV and tuberculosis... 402282fc91 Signs and symptoms of early pregnancy may include missed periods, tender breasts, morning sickness (nausea and vomiting), hunger, implantation bleeding, and frequent urination. Pregnancy may be confirmed with a pregnancy test. Methods of contraception are used to avoid pregnancy. 402282fc91 Drugs taken in pregnancy including over-the counter-medications, prescription medications, nutritional supplements, recreational drugs, and illicit drugs may cause harm to the mother or the unborn child. Tobacco, alcohol, marijuana, and illicit drug use while pregnant may be dangerous for the unborn baby and may lead to severe... 402282fc91 == Early pregnancy == 402282fc91

https://lb1-s245-pub.pressidium.com/=c/text/niche?EPUB=how_much-weight-gain_is_normal_during-pregnancy
[https://lb1-s245-pub.pressidium.com/\\$l/pub/file?PAGE=can-we-drink-milk_after_eating_chicken](https://lb1-s245-pub.pressidium.com/$l/pub/file?PAGE=can-we-drink-milk_after_eating_chicken)
[https://lb1-s245-pub.pressidium.com/\\$c/doc/exe?DOC=how_much_does-a_2-month_old_eat](https://lb1-s245-pub.pressidium.com/$c/doc/exe?DOC=how_much_does-a_2-month_old_eat)
https://lb1-s245-pub.pressidium.com/^h/text/find?PDF=sarna_humana-quanto-tempo_dura
https://lb1-s245-pub.pressidium.com/~h/short/go?PPT=structure_of_seed_diagram
https://lb1-s245-pub.pressidium.com/+o/ref/slug?PDF=what_is_monotonic-preference
https://lb1-s245-pub.pressidium.com/_g/book/link?PLAY=seltzer_water_and_club_soda
https://lb1-s245-pub.pressidium.com/+x/edu/go?KINDLE=c-reactive-protein_blood_test_mean
https://lb1-s245-pub.pressidium.com/=q/ebook/dl?DOC=sangrado_en-el_embarazo_sin_dolor
https://lb1-s245-pub.pressidium.com/_s/pdf/link?PDF=can-fasting_reduce_blood_pressure
https://lb1-s245-pub.pressidium.com/=x/pdf/key?EBOOK=white-spots_in-mouth
https://lb1-s245-pub.pressidium.com/~t/course/url?KINDLE=test_dna-for_free
https://lb1-s245-pub.pressidium.com/!c/ref/mirror?EBOOK=what_does_high_in_calcium-mean
https://lb1-s245-pub.pressidium.com/^z/slide/data?PPT=dreams-why_do-we_dream